

**3613****Topic Category:** 4063-ASIP Inflammation and inflammatory mediators**First Author:** Alexander Wein

Emory University

Department of Microbiology and Immunology

Room 3178 Atlanta, GA 30322

United States

**Phone:**

awein@emory.edu

**First Author is a:** Graduate Student**First Author is a member of:** Not a Member of a Host EB Society**First Author Degree:** BA, BS, or equivalent**Presentation Preference:** Oral**Sponsor:** Asma Nusrat**Sponsor Phone:** (734)764-5712

anusrat@med.umich.edu

**Sponsor's Society:** Pathology - American Society for Investigative Pathology (ASIP) - Host Society**Keywords:** 1. influenza 2. alveolar macrophages

## **IL-36 $\gamma$ Promotes Alveolar Macrophage Survival During Influenza Infection, Limiting Morbidity and Mortality**

Alexander N Wein<sup>1</sup>, Paul Dunbar<sup>1</sup>, Sean R McMaster<sup>1</sup>, Timothy L Denning<sup>2</sup>, Jacob E Kohlmeier<sup>1</sup>. <sup>1</sup>Department of Microbiology and Immunology, Emory University, Atlanta, GA, <sup>2</sup>Institute for Biomedical Sciences, Georgia State University, Atlanta, GA

Influenza viral illness has been a high priority research topic in public health and immune pathogenesis, but the roles of individual cytokines in controlling the infection remains unclear. The IL-36 family of proteins is made up of novel IL-1 family members that have shown key roles in lung immunity to bacterial infections, but the role of IL-36 $\gamma$  in influenza infection is unknown. We investigated the function of IL-36 $\gamma$  during infection with low and high pathogenesis strains of influenza. *Il36g* mRNA is upregulated in the lung following influenza infection with both H1N1 and H3N2 influenza. Genetic deletion of *Il36g* results in greatly increased morbidity, mortality, and viral titers compared to mice replete for the gene. This difference correlates with increased pro-inflammatory cytokines in the airways of mice early during infection. Flow cytometric analysis of the immune compartment revealed a dramatic loss of alveolar macrophages by three days post-infection. Previous studies have shown that alveolar macrophages have a key role in limiting early viral replication and preventing death during lethal influenza infection. We observed that the alveolar macrophages in *Il36g*<sup>-/-</sup> mice have an M2-like phenotype at baseline, possibly contributing to their rapid depletion. Transfer of WT alveolar macrophages to *Il36g*<sup>-/-</sup> mice prior to influenza infection leads to increased survival compared to mock-transferred *Il36g*<sup>-/-</sup> mice. These data indicate that IL-36 $\gamma$  is a key protein in driving alveolar macrophage survival during influenza infection and protecting against severe disease.

### **Support or Funding Information**

Research Support: R01 HL122559